



Stuart Barton, M.D., F.A.C.S.
 Robert N. Gebhart, M.D., F.A.C.S.
 Quinten M. Vanderwerf, M.D., F.A.C.S.
 Christopher Walz, M.D., F.A.C.S.

The following is a partial list of medications that contain ingredients that may interfere with your blood's ability to clot. These medications are **aspirin-containing products, arthritis medications, sleep aids and some herbal products**. Please discontinue these medications 10 – 14 days prior to your surgery. Taking these medications 10 – 14 days of surgery could lead to postponement or cancellation of your surgery. Continue to take your other medications under the direction of your physician. Please check with your prescribing physician, pharmacist or our office should you have any questions.

| | |
|-----------------------|----------------------|
| Advil | Indocin/Indomethacin |
| ALka Seltzer | Meridia |
| Anacin | Midol |
| Anaprox | Nalfon |
| APC | Naprosyn |
| Ascriptin | Norgesic |
| Apergum | Nuprin |
| Aspirin | Nyquil |
| Bayer Aspirins | Oxycodone |
| Bufferin | Pamprin |
| Butazolidin | Percodan |
| Coricidin | Phenaphen |
| Darvon/Darvocet | Propoxyphene |
| Dristan | Robaxin/Robaxis |
| Duragesic | Sine-Aid |
| Ecotrin | Sine-Off |
| Empirin | Sodium Salicylate |
| Equagesic | Sominex |
| Excedrin | St. John's Wart |
| Feldene | Talwin |
| Fenoprofen | Vanquish |
| Fiorinal | |
| Fish Oil | |
| Flax seed Oil | |
| Four-Way Cold Tablets | |
| Ibuprofen | |

Please consult your prescribing physician before discontinuing any following Medications:

- Aggrenox
- Brilinta
- Clopidogrel
- Coumadin/warfarin (generic)
- Dipyridamole
- Heparin
- Persantine
- Plavix
- Pletal
- Pradaxa
- Ticagrelor
- Ticlopidine
- Ticlid
- Trental
- Vimovo
- Xarelto